

10/5/06, 404

PLT 6/21/2003

AD 7/23/02 claim

look up

Priority

8

Claims

5,746,948

1. A method of preparing a membrane having an affinity for biomolecules comprising the steps:

- (a) providing a microporous membrane;
- (b) reacting said membrane of step (a) with a reagent containing a functional group to form a functionalized membrane containing reactive functional groups on the surfaces thereof;
- (c) contacting said functionalized membrane of step (b) with a solution containing an affinity ligand to couple said ligand to said functional group to form a biologically active membrane;
- (d) washing said biologically active membrane with a washing solution containing a volatile organic compound that is miscible with said washing solution; and
- (e) drying said biologically active membrane.

2. The method of claim 1 wherein said functional group of step (b) is an aldehyde.

3. The method of claim 1 wherein said affinity ligand of step (c) is selected from the group consisting of thiophiles; hydrophobes; reversed phase ligands; dyes; low molecular weight charged or non-charged organic molecules; amino acids and analogs thereof; coenzymes, cofactors and analogs thereof; substrates and analogs thereof; endocrine and exocrine substances; enzyme substrates, enzyme inhibitors and analogs thereof; fatty acids, fatty acid derivatives, conjugated fatty acids and analogs thereof; nucleic acids; monomers and analogs and derivatives thereof; polymers and oligopolymers and analogs and derivatives thereof; high molecular weight carbohydrates; glycolic conjugates; proteins and oligomers, subunits and parts thereof; peptides, polypeptides and analogs and derivatives thereof; lectine; antibodies and parts thereof; fusion proteins; haptens; enzymes and subunits and parts thereof; structural proteins; receptors and parts thereof; xenobiotics; pharmaceuticals and pharmaceutically active substances; alkaloids; antibiotics; and biomimicking substances.

Ex 2 + 3

Ex 2 + 3

Col 5 + Col 6

54-56 24-65

17

4+3

9

4. The method of claim 3 wherein said affinity ligand is Protein A. 11

5. The method of claim 1 wherein said washing solution of step (d) is aqueous-based.

5

6. The method of claim 5 wherein said washing solution is a phosphate buffered saline solution.

7. The method of claim 6 wherein said volatile organic compound of step (c) is glycerine.

10

8. The method of claim 1 wherein step (c) is conducted at ambient temperature.

15

9. The method of claim 1 wherein said microporous membrane of step (a) is a polymeric membrane selected from the group consisting of cellulose acetate, cellulose nitrate, polyamide, polyethersulfone, polypropylene and polyvinylidene fluoride.

20

10. The method of claim 9 wherein said microporous membrane has an average pore diameter of from about 0.01 to about 15 microns and a thickness of from about 100 to about 500 microns.

11. The membrane product of the method of any of claims 1-4.

25

12. The membrane product of claim 11 stored in a dry state in a substantially anaerobic atmosphere. Ex4

13. At least one of the membrane product of claim 11 in a filtration housing having a fluid inlet and a fluid outlet wherein said at least one membrane product is situated between said inlet and said outlet.

30

Ed 12

20-25

10

14. The membrane product of claim 13 wherein said filtration housing contains three of said membrane products.

103.2h

112 2nd